





Gases

$$\frac{PV}{T} = k$$



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases- A State of Matter

Properties

- may be compressed
- expand to fill their containers uniformly
- have low densities
- may be mixed
- exert constant uniform pressure on the walls of their containers



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Kinetic (Moving) Theory of Gases



Gases are composed of molecules in constant motion


Gas molecules move in random directions

Molecules of a gas collide frequently with each other & with vessel walls
(why gases mix to uniformity & fill all portions of the containment vessel)

Gas molecules move with an average velocity at a given temperature.
(the average energy of molecules in a gas is the same for all substances)

Distance between gas molecules \gg than size of the individual molecules
(why gases can be compressed)


Molecules are perfectly elastic ... no energy is lost when molecules collide
(If not-elastic, the temperature of a gas mix would always decrease with time)



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Pressure is result of molecular impact on container walls



$$\text{Pressure} = \text{force/area}$$

Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT



General Gas Law




$$\frac{p_1 v_1}{t_1} = \frac{p_2 v_2}{t_2}$$



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Proportional Thinking



$$\frac{p v}{t} = k$$

Variables change to keep k constant

If P constant:

$$\frac{v}{t} = k$$

v and t change
(increase or decrease)
in same direction



If V constant:

$$\frac{p}{t} = k$$

p and t change
(increase or decrease)
in same direction



If T constant:

$$p v = k$$


p and v change
(increase or decrease)
in opposite direction



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases In The News





Chlorofluorocarbons (CFC) Deplete Ozone
Ozone loss increases amount of harmful UV reaching the earth
Skin cancers on the rise in Australia and New Zealand
Loss of all Ozone will eliminate life on the surface of the earth

Copyright Larry P. Taylor, Ph.D. All Rights Reserved


LPT

Gases In The News


Annual Greenhouse Gas Emissions by Sector



Carbon Dioxide
(72% of total)



Methane
(18% of total)





Nitrous Oxide
(9% of total)



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT


Rapid Increase In Greenhouse Gases



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

Gases In The News

There appears to be a correlation:
Greenhouse Gases and Temperature



NYC==> 13" in last century

LPT

Gases In The News



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases In The News


Lighter than air aircraft re-emerging
Best long-range, low cost, heavy transport vessels known



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Helium Filled Balloon As Bridge Support



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Weather Balloons



NOAA facility at White Lake
Balloon filled with H₂
Launched world-wide every day at same time (7 am/pm ± 1 hour)
Lasts ~ one-half hour

Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

High-Tech Balloons



Project Loon
Google Send & Receive WiFi
Neptune
Space Tourism



China
International WiFi Eavesdropping



US Border Patrol
Surveillance Balloon

Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT


Yes, Indeed, Chemistry is a Gas!




Both H_2 and CH_4 can be biologically produced
So, biological “flamers” are conceivable

Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT



Gas Law Lab (Boyle & Charles)



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT


Gases Lab – 50 ml Syringe



Procedure

Part A: Volume – Temperature (Charles)

- Fill a 250-mL Erlenmeyer flask with about 200 mL of distilled water
- Volume measuring device: rubber stopper, 90° glass bend, rubber tubing, & 50 mL syringe:




Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases Lab - Erlenmeyer Flask



- Cap the hole in the rubber stopper with your finger. Drain the syringe through the stopper.
- When ~ 10-15 mL of water remains in the syringe,
- Insert the stopper into the Erlenmeyer flask and invert it → all in the same motion
- Keep the Erlenmeyer flask inverted at all times



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases Lab - 2 L Beaker

- Place the inverted Erlenmeyer flask in a 2000 mL beaker
- Need 2 L beaker to hold the water bath for temperature control



Copyright Larry P. Taylor, Ph.D. All Rights Reserved


LPT

Gases Lab – Ring Stand Ring

- Set the beaker on a ring stand.
- To keep flask from floating, add 2 rubber stoppers on the top of the flask
- Add a wire gauze on top of the rubber stoppers
- Add an iron ring so that it rests on top of the wire gauze.
- Clamp the iron ring to a ring stand



When water is added, the ring/gauze/stoppers hold the flask in place



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases Lab – Ring Stands



- Use 3-fingered clamp on second ring stand to hold syringe in place
- Let the hose between the flask & the syringe hang over the side of the lab bench
The syringe will have to be moved to equilibrate levels



Syringe must be free to move up & down
Control by sliding clamp on ring stand

Add thermometer

This is the “flexible U”

Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases Lab – Recording Volume



Set syringe water level = flask water level

Sight through syringe,
Move syringe to align levels
Record volume to nearest tenth of mL

Most common errors:
Not setting equal water levels
Not equilibrating temperatures

Record initial reading
This will be initial for all measurements



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases Lab – Volume vs. Temperature

- Heat water (hotplate) until at least 60 °C; carefully pour into 2 L beaker
- The flask in the 2 L beaker must be completely covered by the hot water
- Add ice to cool the bath to ~50 °C
- **RECORD the VOLUME as the SYRINGE READING for the T = 50.0 °C**
- Add ice, with constant stirring, to lower the temperature
- Record Temperature & Syringe Volume for T = 40.0, 30.0, 20.0, and 10.0 °C.


At each step:
Must move syringe up and down
So level of syringe = flask



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases Lab – What's Happening



As Gas Volume (Trapped Gas) changes,

The change is measured here
(Data point – initial reading)

Total Volume = Trapped Gas + Δ Syringe



Copyright Larry P. Taylor, Ph.D. All Rights Reserved


LPT

Gases Lab – Vacuum Gauge



Procedure

Part B: Volume – Pressure (Boyle)



Vacuum Gauge (mmHg below atmospheric)

Drierite (Desiccant-keeps water out of gauge)

Safety Flask (keeps water out of vacuum Line)

Needle Valve (controls vacuum) by air bleed



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases Lab – T-Valve



Needle Valve – Bleeding Air Here Controls Vacuum



To vacuum



To syringe



Use vacuum tubing:

Short Tubing Connects to the syringe

Long Tubing Connects to the Vacuum Line

Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases Lab – Complete Apparatus

- Completely open the needle valve
 - Carefully open the needle valve
 - Watch fluid level
(too much vacuum → liquid moves into vacuum line)
- All adjustments should be done with the needle valve
Close the needle valve to obtain a reading of 50 mm Hg
- Adjust the syringe water level = water level in the flask
 - RECORD the VOLUME of SYRINGE READING
 - Repeat for 100, 150, and 200 mm of Hg
 - Read the barometer (same reading for all measures)



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Gases Lab – Barometer


Behind Instructor's Desk

Read Atmospheric Pressure in mmHg

Pressure in the Apparatus is:

Atmospheric (barometric) Pressure – Vacuum Gauge Reading

Instructor will write pressure on the blackboard



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

What's Happening



The volume of trapped air (in the flask) is changing with pressure
This change is measured with the syringe

Total gas volume (for each different pressure point):

Trapped air (measured later) plus syringe volume change

Barometric pressure is the pressure of the air that surrounds the apparatus

The vacuum gauge indicates the decrease in pressure within the apparatus

Pressure on the gas = barometric pressure – vacuum gauge pressure.



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT


Gases Lab – Measuring Trapped Gas Volume

Disassemble the apparatus

Empty the flask to the sharpie mark

Measure volume with a 50 mL graduated cylinder

Record this as the Trapped Air (Gas) Volume.



Fill to mark

Pour into 50 mL graduate

Record Volume

Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Data Interpretation

Plot temperature versus volume graph for Part A of this experiment.

(Temperature values on the x-axis; higher temperatures on the right)

Plot the corresponding total volume of the gas in the flask values on the y-axis

Conclusion

Indicate:

Change of gas volume with change in temperature

Product of the pressure and volume of the gas at constant temperature



Hints:

If $V/T = \text{a constant}$ (within experimental error), then V & T are directly proportional

If $P \times V = \text{a constant}$ (within experimental error), then P & V are inversely proportional

Questions:

“Molecular explanation” means think about what are the molecules doing ... do not just recite the gas laws. (i.e. use Kinetic Theory of Gases)



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

This Lab Always Produces Good Results



Somewhere, Charles & Boyle Are Smiling



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT

Let's Boldly Go Explore Today's Lab



Copyright Larry P. Taylor, Ph.D. All Rights Reserved

LPT